



# How to taper fluvoxamine safely

A clear, practical walk-through of how people reduce fluvoxamine gradually — written to read alongside the advice your prescriber gives you, never instead of it. About a 7-minute read.

⚠️ **This is a dose conversion calculator, not a tapering plan.** Get a safe tapering plan from your healthcare team before starting any reduction. The [Maudsley Deprescribing Guidelines](#) and the [RELEASE Toolkit](#) are good resources to discuss with your doctor, or [book a telehealth appointment with the TaperMate Clinic](#) to get a personalised tapering plan.

## Why come off slowly

Stopping fluvoxamine is usually harder than starting it. The relationship between the dose you take and its effect on the brain isn't a straight line — at lower doses, even a small reduction in milligrams can be a large change in effect. This is why prescribers increasingly recommend **hyperbolic tapering**: smaller and smaller reductions as the dose gets lower, rather than fixed steps.

In practice that means the standard strengths — designed for the treatment range, not for coming off — often can't give you the small doses the end of a taper needs. That's where a method like the one below, so you can measure a precise fraction, comes in.

## The liquid method, step by step

1. Cut a tablet to the portion you need with a pill cutter.
2. Crush the piece between two spoons over a small plate.
3. Stir it into a measured volume of water to make a known concentration — it will look cloudy, which is expected.
4. Stir again immediately before drawing your dose; the active ingredient settles quickly.
5. Measure your prescribed dose with an oral syringe and take it.
6. Make a fresh liquid each day and discard any unused liquid in the rubbish.

[Open the fluvoxamine calculator to work out your dose →](#)

# Your step-by-step taper schedule

This is the Fluvoxamine schedule from the RELEASE Toolkit, reproduced with permission. It's a starting point to **discuss with your prescriber** — you can pause, slow down or speed up depending on how you feel. Aim to reduce roughly every 2–4 weeks.

STEP	DAILY DOSE	LIQUID OPTION (2MG/ML)	CAPSULES OPTION
1	100mg	2 × 50mg tablets	—
2	75mg	1½ × 50mg tablets	—
3	50mg	1 × 50mg tablet	—
4	35mg	½ × 50mg tablet & 5ml liquid	½ × 50mg tablet & 2 × 5mg capsules
5	25mg	½ × 50mg tablet	—
6	15mg	7.5ml liquid	3 × 5mg capsules
7	12mg	6ml liquid	2 × 5mg & 1 × 2mg capsules
8	10mg	5ml liquid	2 × 5mg capsules
9	8mg	4ml liquid	4 × 2mg capsules
10	6mg	3ml liquid	3 × 2mg capsules
11	4mg	2ml liquid	2 × 2mg capsules
12	2mg	1ml liquid	1 × 2mg capsule
13	Stop	You've completed the taper 🎉	

Highlighted steps are the most important — do not skip them.

- Do **not** skip the final small-dose steps — they're the most important for preventing withdrawal.
- Don't skip days, alternate days, or suddenly stop.
- If withdrawal symptoms appear, you can return to your previous dose; when ready, reduce more slowly.

Schedule © The University of Queensland (RELEASE Toolkit), reproduced with permission. Dosing guidance: Dr Mark Horowitz.

# Measuring smaller doses accurately

Use the right tool for your method – an **oral syringe** for liquids (your pharmacist can supply 1 mL, 5 mL and 10 mL sizes), or a **milligram scale** for weighing. Choose the smallest measuring tool that fits your dose: it's far more accurate for tiny amounts.

- Measure slowly and double-check before you take your dose.
- Read at eye level, against the syringe plunger's flat edge.
- Keep the concentration or method the same each day so doses stay consistent.

# What withdrawal can feel like

Withdrawal effects vary a lot between people. They often come in **waves** – harder days followed by **windows** of feeling more like yourself. Common, usually-manageable effects include dizziness, "brain zaps", nausea, vivid dreams, irritability and low mood. They tend to ease if you hold at your current dose for a while before reducing again.

**Seek advice promptly** if you have thoughts of harming yourself, severe or worsening mood, or symptoms that don't settle. Contact your prescriber, or in an emergency call **000**. In Australia you can also call **Lifeline on 13 11 14**.

# Questions for your appointment

- How quickly is it safe for me to reduce, and by how much each step?
- How long should I hold at each dose before the next reduction?
- What should I do on a bad day – hold, slow down, or pause?
- Which symptoms mean I should contact you sooner?
- Would a compounded oral liquid be more accurate for my lowest doses?

NEXT STEP

## Plan your full taper in TaperMate

This calculator handles one dose. The TaperMate app calculates a full reduction schedule with hold periods, microtapering and symptom monitoring – so each new dose is one tap away, not a daily maths problem.



Download on the  
App Store



Get it on  
Google Play

---

## Sources to discuss with your prescriber

- Horowitz MA, Taylor D. *The Maudsley Deprescribing Guidelines: Antidepressants, Benzodiazepines, Gabapentinoids and Z-drugs*. Wiley, 2024.
- **RELEASE Toolkit** – Reducing & Eliminating LongtErm AntidepreSsant usE. [releasetoolkit.com.au](https://releasetoolkit.com.au)
- **Therapeutic Goods Administration**. Therapeutic Goods (Excluded Goods) Determination 2018, item 14L.